

Evidence of microbially accelerated weathering from a laboratory mesocosm experiment with sequential selective dissolution

Corey R. Lawrence¹, Harun Niron², Tania Timmermann¹, Philip D. Weyman¹, Yun-Ya Yang¹, Daniel
Dores¹, Gonzalo A. Fuenzalida-Meriz¹

¹Andes Ag, Inc., Alameda, California, USA

²Biobased Sustainability Engineering (SUSTAIN), Dept. of Bioscience Engineering, Univ. of Antwerp, Antwerp, Belgium

Correspondence to: Gonzalo A. Fuenzalida-Meriz (gonzalofuenzalida@gmail.com)

Abstract. Microbially accelerated weathering (MAW) is a promising soil-based carbon dioxide removal (CDR) strategy that leverages beneficial soil microbes to increase weathering of preexisting silicate minerals. This approach does not require addition of a mineral feedstock, greatly reducing the carbon footprint from mining, grinding, transporting, and applying the mineral to land compared with enhanced weathering. A key obstacle to measurement, reporting, and verification for MAW is ensuring that increases in weathering products, such as base cations, are sourced from silicate dissolution rather than redistribution of pre-existing cations from the exchangeable, oxidizable, or reducible soil pools. To address this, we conducted a 63-day mesocosm study with soybean, utilizing soil sequential extractions to track the buildup and distribution of weathering products in soil columns inoculated with *Bacillus subtilis* strain MP1. Our results indicated that MP1-treated soils yield a net increase in available base cations, corresponding to a 4.5% increase in base cation charge relative to control soils. The observed increase in base cations is partitioned between the carbonate and exchangeable soil pools. We estimate that 37% to 67% of the weathering derived base cations formed carbonates, corresponding to a CDR of 0.20 to 0.36 g CO₂ kg⁻¹ soil. These findings suggest that *Bacillus subtilis* MP1 couples silicate mineral dissolution with carbonate precipitation, providing support for MAW as a viable and scalable CDR strategy.

1 Introduction

In addition to reducing carbon emissions, technologies enabling carbon dioxide removal (CDR) are needed to meet the requirements of The Paris Agreement and lessen the impact of anthropogenic greenhouse gas (GHG) emissions (Johnstone et al., 2026; Smith et al., 2024). Soil-based CDR methods, such as enhanced weathering (EW) and microbially accelerated weathering (MAW), are appealing because they can be applied in agricultural systems, resulting in co-benefits for soil health and productivity, and where they can be efficiently scaled. The goal of EW is to increase the dissolution of silicate minerals, which is a natural but slow process that acts as a regulator of GHGs over millennial timescales (Berner, 1997; Walker et al., 1981). There are, however, economic and biogeochemical constraints on this approach that limit its effectiveness in some soil environments. For example, most EW approaches work best in acidic soils, and CDR must be high enough to offset emissions associated with the mining, grinding, transport, and spreading of the feedstock material. Furthermore, the measurement,

reporting, and verification (MRV) required for issuance of high-quality carbon credits from soils can be difficult to efficiently apply at scale. In the case of EW, this has led to a variety of MRV approaches, each with unique costs and benefits (see Clarkson et al., 2024). It is becoming increasingly apparent that there is no ‘one size fits all’ approach. Technological innovation coupled with applied research will continue to be essential for achieving CDR at the scale required to combat climate change.

Weathering of silicate minerals contained in rocks and soil is a natural negative-feedback to increasing atmospheric CO₂, whereby carbonic acid formed from CO₂ and water supplies the acidity needed to drive mineral dissolution. As silicates dissolve, base-cations contained within the minerals are released to solution, and carbon from carbonic acid is converted to carbonate alkalinity (i.e., $\text{HCO}_3^- + \text{CO}_3^{2-}$). In natural settings, these reactions are assumed to be too slow to effectively offset anthropogenic GHG emissions over human timescales. EW approaches increase weathering rates by adding finely crushed mineral feedstocks to soils, accelerating effective weathering rates by increasing the mineral surface area. However, it is becoming increasingly clear that mineral weathering is not strictly an abiotic process; microbes, including bacteria and fungi, have been shown to facilitate and enhance rates of mineral weathering (Banfield et al., 1999; Corbett et al., 2025; Finlay et al., 2020; Timmermann et al., 2025; Verbruggen et al., 2021; Vicca et al., 2022; Wild et al., 2022). There are several hypothesized mechanisms through which microbes may increase mineral weathering, including the production of carbonic anhydrase enzymes that enhance the solubility of CO₂ and formation of carbonic acid (Timmermann et al., 2025; Vicca et al., 2022; Xiao et al., 2015) or through exudation of organic acids that supply acidity for weathering and act as ligands that remove weathering products (Niron et al., 2025; Ribeiro et al., 2020; Vicca et al., 2022). Harnessing this biological potential may provide another valuable tool to accelerate and scale CDR.

Microbe-based approaches for accelerating silicate weathering, such as MAW, can be applied to soil systems without the addition of exogenous mineral feedstocks (Timmermann et al., 2025). By increasing weathering of preexisting soil silicate minerals, upfront costs and carbon emissions can be reduced, allowing for a more efficient and cost-effective CDR process. However, this approach requires soils meeting specific geochemical conditions, and soils must be (1) rich in cation-bearing silicate minerals to be weathered, (2) neutral to alkaline pH (discussed in Yang et al., 2026), and (3) able to precipitate products of mineral dissolution as secondary carbonates. Timmermann et al. (2025) proposed that accumulation of secondary carbonate minerals, facilitated by microbial amendments, serves as an effective proxy for weathering, allowing for MRV approaches based on carbon rather than base cations. When carbonate formation is coupled with the dissolution of silicates containing divalent cations (Mg^{2+} and Ca^{2+}), which consume two moles of CO₂ for every mole of mineral weathered, half of the CO₂ removed is returned to solution. Carbonate formation therefore reduces the efficiency of CDR from silicate weathering, which is why it is undesirable in traditional EW systems. Yet, without the need for mining, crushing, and transporting feedstock and the associated CO₂ emissions, this process can still allow for cost-effective CDR in a system where only microbial amendments are added to a soil. However, one potential concern with such an approach is that increases in secondary carbonate

accumulation could conceivably occur without corresponding increases in silicate weathering, for example, if the divalent cations precipitated in carbonates are sourced from somewhere else in the soil system (e.g., the exchangeable complex). To address this concern and definitively link microbial amendments with increases in silicate dissolution, we conducted an experiment leveraging sequential selective soil extraction, a technique allowing for a comprehensive characterization of the soil weathering system, including tracking of all potential sinks for weathering products (e.g., Steinwider et al., 2026).

In this study, we describe the results of a laboratory mesocosm experiment, where soybeans were grown with and without the addition of a microbial amendment, *Bacillus subtilis* MP1. We have previously demonstrated that, in both laboratory and field experiments, when soils are amended with strain MP1 or similar microbes, the microbial acceleration of silicate mineral dissolution can facilitate CDR (Niron et al., 2024; Timmermann et al., 2025). Additionally, we have shown the potential for these microbial amendments to improve the effectiveness of EW when microbes are added in combination with silicate feedstocks (Yang et al., 2026). However, in both microbial amended and traditional EW systems, it is important to track the fate of weathering products, including base-cations and carbonate alkalinity. Here we used a sequential selective soil extraction procedure (e.g., Niron et al., 2024; Steinwider et al., 2025; Vienne et al., 2026) to trace how the microbial amendment MP1 altered silicate weathering and changed various soil base-cation pools. Data from selective sequential extractions provide a powerful way to track products of silicate dissolution and the potential sinks of weathering products. By tracking all inputs and reservoirs of base cations in the experimental system, along with inorganic carbon and other diagnostic soil properties, this work clearly demonstrates that the microbial amendment MP1 facilitates carbonate accumulation by increasing the dissolution of silicate minerals.

2 Materials and Methods

2.1 Soil mesocosm experiment

We conducted a 63-day soil mesocosm experiment with 12 soil columns to assess the impact of the microbial amendment *Bacillus subtilis* strain MP1, on soil silicate weathering under soybean (*Glycine max*) cultivation. The soil used for the mesocosm experiment, hereafter referred to as SBX70, was a silty clay soil collected from an agricultural field in Stutsman County, North Dakota, USA. Prior to the mesocosm preparation, the field-moist SBX70 soil was sieved through a 10 mm mesh to remove large stones and plant debris, and then manually homogenized. Each mesocosm (35 cm height and 10 cm diameter) was filled with 3 kg of the prepared SBX70 soil to a total height of 30 cm, leaving 5 cm of headspace. After correcting for the moisture content of the soil, this equated to a dry mass of 2.18 kg within each mesocosm.

The experimental design consisted of two treatment groups, each with six biological replicates: (i) untreated control (“UTC”) and (ii) MP1 inoculated (“MP1”) (Figure S1). MP1 is a naturally occurring *B. subtilis* strain isolated from corn roots and rhizosphere soils. Detailed characterization of MP1 can be found in Timmermann et al. (2025). Each mesocosm was planted

with a single soybean seed (Asgrow AG19XF3). To prepare the MP1 treatment, each seed was inoculated with 3.9×10^6 spores of MP1 suspended in 1 mL of distilled water. The concentration of MP1 spores added per mesocosm equates to 3.9×10^3 spores per gram of soil. In the UTC treatment, 1 mL of distilled water per seed was applied instead. The experiment was conducted in a growth chamber at 22 °C (± 5 °C) with a relative humidity of 65% ($\pm 5\%$) and a 16-hour photoperiod. Plants were watered with deionized (DI) water three times a week for the first six weeks.

2.2 Simulated rainfall events and sample collection

Six weeks (42 days) after sowing and MP1 inoculation, six simulated rainfall events were applied to the mesocosms over a two-week period. The total rainfall amount was 1.75 L of DI water per mesocosm, distributed across the six events, with each mesocosm receiving 292 mL DI water per event. The total volume of water was determined as described in Timmermann et al. (2025). The simulated rainfall schedule followed a pattern of two consecutive days of water additions, followed by two days without, repeating three times, for a total of six simulated rainfall events. Leachate was collected from each mesocosm during each of the six discrete rainfall events, thereby accounting for the total leachate flux throughout the experiment. The volume of leachate collected during each wetting event ranged from 0 to 175 ml, with total volumes collected overall all events of 448 ± 16 ml for MP1 amended columns and 464 ± 19 ml for control columns. Following the simulated rainfall period, plants continued to grow under controlled conditions until the end of the experiment on day 63.

After the 63-day experiment, all 12 mesocosms were harvested and soils were divided into three depth intervals for analyses: 0–10 cm, 10–20 cm, and 20–30 cm (Figure S1). A total of 36 soil samples (two treatments with six replicates and three depths per replicate) were collected and homogenized. Then, the samples were air dried, sieved to 2-mm, and ground for soil physicochemical analyses. Soybean plants were harvested separately for above- and belowground biomass assessment.

2.3 Soil mineralogy and chemistry

Soil mineralogy was estimated from combination of the scanning electron microscope (SEM) observations on a Zeiss EVO-10 Variable Vacuum SEM and XRD analysis with the PANalytical X'Pert Pro diffractometer at the Department of Earth and Planetary Science at the University of California (Berkeley, CA, USA). The total elemental composition of the SBX70 soil prior to the onset of the experiment was measured via x-ray fluorescence (XRF) at the GeoAnalytical Laboratory at Washington State University (Pullman, WA, USA), and soil organic matter content was determined by the loss-on-ignition method at 360 °C (Schulte and Hopkins, 1996) to allow for normalization of the elemental composition on a volatile-free basis. Technical details and principles underlying these methods are described in Johnson et al. (1999) and Kelly et al. (1998).

Soil inorganic carbon (SIC) was quantified using a gas chromatography (GC) method described in Yip et al. (2025). Soil bicarbonate (HCO_3^-) and carbonate (CO_3^{2-}) ions were quantified using a saturated paste extract followed by titration (Richards et al., 1954; Yang et al., 2026). Due to the complexity of this measurement, the saturated paste extract was made on a single

depth-composited sample from each column. In this study, “carbonate alkalinity” is defined as the combined concentration of HCO_3^- and CO_3^{2-} ions in the soil.

135 Additional soil physicochemical parameters were analyzed by an external commercial laboratory (Agvise Laboratories, ND, USA). Soil pH was measured using the standard 1:1 soil-to-water ratio method with a calibrated pH meter (Peters et al., 2012). Soil total carbon (TC) was quantified via dry combustion using a Vario MACRO cube elemental analyzer (Elementar Americas Inc., NY), while total soil organic carbon (SOC) was calculated as the difference between TC and SIC (Nelson and Sommers, 1996). Total soil cation exchange capacity was measured using the ammonium acetate displacement method (Sumner and
140 Miller, 2020) and was compared against the sum of exchangeable base cations (described below) to estimate base saturation of the exchange complex, where base saturation (%) is equal to the sum of exchangeable base cations divided by the total cation exchange capacity (both in units of mEq/100g soil).

2.4 Sequential cation extractions

145 Four soil cation pools were sequentially extracted from soil samples to assess changes in cation mobility and distribution: (1) exchangeable cations, (2) carbonate-bound cations, (3) reducible cations, and (4) the oxidizable fraction. Methodology for sequential extractions was adapted from Tessier et al. (1979) and Uhlig and Von Blackenburg (2019), and is detailed in Steinwigger et al. (2025). Our focus was to assess differences between soil and plant systems with and without the MP1 amendment. We did not perform the sequential extraction on the initial soil, homogenized soil, that was used in the experiment.

150

In the first step (exchangeable cations), 10 mL of 1 M NH_4OAc was added to 1 g of air-dried soil, shaken for 1 h, centrifuged, and the supernatant collected. For the second step (carbonates), 5 mL of 1 M CH_3COOH was added and shaken for 2 h; then 1 mL of 3 M NH_4OAc was added. Samples were brought to 10 mL with DI water, centrifuged, and supernatants collected. In the third step (reducible — oxide/hydroxide bound cations), 5 mL of 0.05 M NH_2OH in 1 M HCl was added. Samples were
155 heated at 80 °C for 5 h with manual shaking every 30 min, then treated with 3 M NH_4OAc , diluted to 10 mL, centrifuged, and collected. In the final step (oxidizable — organic-bound cations), 4 mL of 30% H_2O_2 in 0.01M HNO_3 was added, and samples were heated at 70 °C with periodic shaking. After 2 h, 3 mL more H_2O_2 was added, and heating continued for 3 h. Then 1 mL of 3 M NH_4OAc was added, samples were diluted to 10 mL, centrifuged, and collected. Between each step, the soils were rinsed with 10 mL of DI water. All centrifugations were done at 4000 rpm for 10 min, and all extracts were filtered through
160 0.45 μm filters.

Elemental concentrations were measured using ICP-OES (iCAP 6300 Duo, Thermo Scientific) for Ca, Mg, K, and Na in all cation pools, and for Al, Fe, and Si in the last two pools. Samples were acidified (19:1, 2% HNO_3 :sample; TraceMetal Grade, Fisher Chemical) prior to analysis. Calibration used a multi-element standard (CPAChem), with yttrium (Merck) as the internal

165 standard. All standards were diluted in the respective extraction media to ensure matrix-matched calibration and sample conditions.

2.5 Plant biomass & chemistry

At the end of the experiment, above and belowground plant biomass was harvested from each column. Samples were oven
170 dried at 60 °C, then weighed prior to grinding for chemical analyses. Aboveground dry biomass samples were analyzed for elemental composition by ICP-OES following acid digestion at Agvise Laboratories (Northwood ND, USA). Due to lower sample mass from individual columns, an equal mass of belowground biomass from replicate columns were combined, yielding two pooled samples per treatment. The belowground biomass samples were analyzed by ICP-OES following acid digestion at The Pennsylvania State University Agricultural Analytical Service Laboratory (State College PA, USA). To calculate the total
175 cation content attributable to plant above and belowground biomass, measured cation concentrations in plant tissue (mg/kg) were multiplied by plant tissue dry weight.

2.6 Leachate chemistry

Leachate samples, collected from mesocosms following each simulated rainfall event, were analyzed for several
180 biogeochemical parameters. Leachate pH was measured with a benchtop probe following a multipoint calibration. Bicarbonate and CO_3^{2-} ions were quantified via titration with 0.0125 M H_2SO_4 with phenolphthalein and methyl orange indicators (Richards et al., 1954). Dissolved inorganic carbon (DIC) in leachate was measured on 4 mL of sample following the GC method described in the previous section. Remaining leachate samples were stored frozen and submitted to Agvise Laboratories (Northwood, ND, USA) for analysis of concentrations of soluble cations Ca^{2+} , Mg^{2+} , K^+ , and Na^+ (APHA, 1998) as well as
185 major anions including NO_3^- , SO_4^{2-} , and Cl^- . Soil leachate data was assessed for charge balance. Positive charge offsets were assumed to result from sample re-equilibration during the collection phase and were corrected by adjusting total carbonate alkalinity ($\text{HCO}_3^- + \text{CO}_3^{2-}$) by the corresponding magnitude of the offset (Tosca and Tutolo, 2023).

2.7 Statistical analyses

190 Based on the distribution and homoscedasticity of the data, either a parametric Student's t-test (normal distribution and equal variance) or a nonparametric Mann–Whitney U test (Wilcoxon rank-sum) was applied to evaluate significant differences ($p < 0.05$) in soil properties and leachate chemistry among treatments. Statistical analyses of soil and water physicochemical properties were performed using the GraphPad Prism version 10 (GraphPad Software, Boston, Massachusetts, USA) or using tools from the tidyverse packages (Wickham et al., 2019) in R (v4.5.0, R Core Team, 2021).

195 3 Results

3.1 Bulk soil and leachate chemistry

3.1.1 Initial soil mineralogy and elemental chemistry

The SBX70 soil used for this experiment contained roughly 65% quartz and 28% feldspars, including albite, oligoclase, and orthoclase is roughly equal amounts (Table S1). The corresponding elemental composition constrains the overall abundances of Ca, Mg, K, and Na in the solid phase at the onset of the experiment (Table S2).

3.1.2 Soil and leachate pH

The initial pH of the SBX70 soil was 7.1 ± 0.03 ($N = 3$). The volume weighted pH of column leachate exhibited a small but significant increase ($p < 0.05$, $N = 6$) in the MP1 treated (8.44 ± 0.04) compared to control soils (8.30 ± 0.06) (Figure 1A). At the end of the experiment, the mean value of soil pH was higher for MP1-treated columns compared with UTC, but the difference was not statistically significant (Figure 1B).




Figure 1: Leachate pH (A) and soil pH (B) in MP1-treated and untreated control (UTC) columns. Wilcoxon signed-rank test, * $p < 0.05$, ns: not significant, $N = 6$. Note that y-axes are scaled to the data range to clearly visualize the distribution and variability across treatments.

3.1.3 Alkalinity and inorganic carbon in soil and leachate

At the end of the 63-day experiment, soil carbonate alkalinity ($\text{CO}_3^{2-} + \text{HCO}_3^-$), measured from soil titrations, was significantly higher ($p < 0.05$, $N = 6$) in MP1 compared with UTC columns (Figure 2A). Alkalinity in leachate is reported as the total cumulative flux normalized by the mass of soil in each column (mEq/kg soil) so that it can be directly compared to soil measurements. Leachate alkalinity was similar between treatments, thus, differences between treatments in the total alkalinity of the system (soil + leachate), were mainly driven by the solid phase, resulting in a higher total alkalinity for MP1-treated soils in comparison to UTC soils (Figure 2A). Inorganic carbon in leachate and soil showed similar trends, with higher mean values of SIC and TIC in the MP1 treated columns ($\approx 37\%$ increase over UTC) (Figure 2B).




Figure 2: Carbonate alkalinity (A) and inorganic carbon (B) in MP1-treated and untreated control (UTC) columns. Student's t-test, * $p < 0.05$, ns: not significant, $N = 6$.

3.1.4 Soil total, inorganic and organic carbon

Total soil organic carbon (SOC) from the MP1-treated soil ($2.67 \pm 0.03\%$) was significantly higher than UTC ($2.58 \pm 0.02\%$) in the 20-30 cm soil increment ($p < 0.05$, $N = 6$) (Figure 3A). Total inorganic carbon (TIC) concentrations were near the detection limit of our method for all soils in this study, and though not significant, mean values of MP1-treated soils were consistently higher in TIC across all depth increments (Figure 3B), and at the column level (Figure S2B). When considering the entire column, total carbon was significantly higher ($p < 0.05$, $N = 6$) in the MP1 treatment versus UTC (Figure 3C), and with depth resolution, significant increases in the MP1-treated soils were observed in the 20-30 cm increment (Figure S2A).

3.1.5 Soil cation exchange capacity and columns base cation fluxes

Values of total cation exchange capacity ranged from roughly 40 to 47 mEq/100g soil across all treatments and depths, and no clear patterns or significant differences were observed (Figure S3). The total flux of cations in column leachate (calculated as leachate volume x concentration) was 5% higher in the MP1 treatment, and all individual base cations were higher in MP1 compared to UTC samples, but differences were not statistically significant (Figure S4).




Figure 3: Total organic carbon (A), total inorganic carbon (B) and total carbon (C) in MP1-treated and untreated control (UTC) columns. Wilcoxon signed-rank test, * $p < 0.05$, ns: not significant, $N = 6$. Note that y-axes are scaled to the data range to clearly visualize the distribution and variability across treatments.

3.1.6 Plant cation content

Plant biomass was slightly elevated in UTC compared with MP1 columns, but the concentration of base cations tended to be higher in plants from the MP1 column. Neither trend was statistically significant (data not shown), and the overall mass of individual base cations attributable to plant matter was similar for both UTC and MP1 treatments. There were no significant

differences observed for the total mass of Mg and K between treatments. The total mass of Ca was higher in MP1 treatment compared with UTC ($p < 0.1$); however, Na was significantly lower in the MP1 treatment compared with UTC ($p < 0.05$, Figure S9).

245 **3.2 Sequential extractions results**

3.2.1 Exchangeable fraction

There were no significant differences in exchangeable base cations between treatments, though mean values of exchangeable Ca were consistently higher in MP1-treated columns compared with UTC across all soil depths (Figure S5). Differences in base saturation, calculated as the ratio of the sum of exchangeable base cations to the total cation exchange capacity, were not

250 statistically significant but the mean values were consistently higher for MP1-treated versus UTC columns (Figure S6).




Figure 4: Concentration per depth increment of base cations in the carbonate pool in MP1-treated and untreated control (UTC) columns. Student's t-test, * $p < 0.05$, ns: not significant, $N = 6$.

3.2.2 Carbonate fraction

255 The carbonate extraction yielded significant differences between MP1 and UTC treatments. In the 10 to 20 cm depth increment, all base cations (Ca, Mg, K, and Na) were significantly higher in the MP1 treatment (Figure 4). Concentrations of K were also significantly greater in the 0-10 and 20-30 cm depth increments.

3.2.3 Reducible fraction

260 Except for Na in the 10-20 cm depth increment, which was significantly higher in the MP1-treated columns, masses of elements extracted from the reducible pool were comparable between the MP1 and UTC treatments of the top two depth increments (Figure S7). In contrast, masses of Mg and Si extracted from the reducible pool of the 20-30 cm depth increment were significantly greater in the MP1-treated columns compared with UTC (Figure 5). The mean values of all other elements measured (Al, Fe, Ca, K, and Na) were also higher in the MP1-treated columns but differences were not significant at the $p < 0.05$ threshold. The normalized molar ratio of Fe:Mg:Si found in the reducible fraction of both treatments was on the order of

265 1:2.5:4 consistent with Si-Mg rich Fe-(hydr)oxides.




Figure 5: Concentration at the bottom depth of base cations in the reducible pool in MP1-treated and untreated control (UTC) columns. Student's t-test, * $p < 0.05$, ns: not significant, $N = 6$.

3.2.4 Oxidizable fraction

270 With one exception, there were no significant differences in elemental masses extracted from the oxidizable pool of the MP1 and UTC columns (Figure S8). The exception was Na in the 10 to 20 cm increment, which was near the detection limit for all MP1 columns.

3.3 Cation mass-balance

At the end of the 63-day experiment, the total mass of base cations recovered across all biogeochemical pools measured in this study was higher in the MP1-treated soils compared with UTC soils, for Ca, Mg, and K (Figure 6). In contrast, the total amount of Na recovered was higher in the UTC soils. For Ca and Mg, the two dominant cations in the system, the exchangeable pool represented the largest proportion of the total recovered mass, followed by the carbonate pool (for Ca) and reducible pool (for Mg). Only a small fraction of these cations was associated with the oxidizable, plant biomass, or leachate pools. The pattern for K was consistent with those observed for Mg: most of the recovered K resided in the exchangeable and reducible pools. Whereas the pattern for Na was consistent with those observed for Ca: the two dominant pools were the exchangeable and the carbonate pools. Contrary to the other three cations, the amount of Na recovered from the reducible pool was negligible (Table S3).




Figure 6: Cation mass-balance. Total mass of cations (in mg) per column, and their distribution between measured pools at the end of the 63-day experimental period. The carbonate, exchangeable, reducible, and oxidizable fractions are based on selective sequential extraction measurements.

4 Discussion

4.1 A conceptual framework for microbe-mediated weathering

Leveraging the soil microbiome to promote carbon dioxide removal represents a promising approach to help offset anthropogenic carbon emissions (Fierer and Walsh, 2023). One such strategy involves using microbes to increase silicate mineral weathering, which converts atmospheric CO₂ to HCO₃⁻ and carbonates. For example, previous experiments using the plant growth-promoting rhizobacterium *B. subtilis* MP1 in combination with plants (Timmermann et al., 2025; Yang et al., 2026) or other *B. subtilis* strains have suggested promising increases in mineral weathering rates in both laboratory and field environments (Corbett et al., 2025; Hopf et al., 2009; Niron et al., 2024; Song et al., 2007). This highlights the potential use of microbial amendments as a CDR strategy, and it is hypothesized that increases in mineral weathering driven by these microbes support plant growth through acquisition of rock-derived nutrients (e.g., P, Ca, Mg, K, Fe), which are essential for plant growth and development (Dong et al., 2022; Samuels et al., 2020). Importantly, we conducted this experiment in the presence of plants (*Glycine max*) because microbial amendments require a consistent source of carbon (e.g., root exudates) to thrive and root respiration concentrates and redirects atmospheric CO₂ belowground, which is necessary to support weathering via the mechanisms proposed. In other words, the impact of microbial amendment is dependent on the presence of a suitable plant host.

Timmermann et al. (2025) presented a conceptual framework for how these microbes may drive silicate mineral weathering, in collaboration with the plant host. In this framework, microbes facilitate the production of acidity (H⁺) via carbonic anhydrases driven catalysis of the conversion of plant-respired CO₂ to HCO₃⁻ (Supuran, 2016), and subsequent precipitation of carbonate (which also generates H⁺). *Bacillus* species are generally understood to influence soil pH (Wang et al., 2025), and the biologically generated acidity is assumed to support further proton-promoted weathering, while the products of weathering are removed from solution or mobilized away from the mineral to facilitate continued dissolution (Figure 4 in Timmermann et al., 2025). This conceptual framework has support in the literature (McConnaughey and Whelan, 1997) but has yet to be fully validated in the context of microbially accelerated weathering. In particular, questions remain regarding the spatial configuration of these processes. However, at bulk soil scale, experimental evidence provides support for the role of secondary carbonate formation in the context of MP1 amendment (Timmermann et al., 2025). Nonetheless, one concern is that the observed increases in soil carbonate associated with the microbial amendment may source divalent cations from somewhere other than the silicate weathering. To address this question, we conducted a soil mesocosm experiment following the approach described in Timmermann et al. (2025) and Yang et al. (2026), this time incorporating a sequential selective extraction procedure (Niron et al., 2024; Steinwider et al., 2025, 2026; Vienne et al., 2026) to trace base cations in available soil reservoirs.

4.2 Bulk soil response to MP1 amendment

Despite being well suited for the MAW approach, with roughly 28% (v/v) feldspars and an initial soil pH of 7.13, the impact of the MP1 microbe amendment in this 63-day long experiment was not entirely apparent from bulk soil and leachate data. The flux-weighted pH of the soil leachate was significantly higher from the MP1 columns (Figure 1A), implying a greater consumption of protons, possibly through increased weathering. Soil pH was also elevated in the MP1 treatment compared with control, though differences were not significant (Figure 1B). Similarly, carbonate alkalinity and inorganic carbon content were each higher in the MP1-treated columns (Figure 2), but only soil alkalinity was significant. There were no significant differences in the leachate flux of alkalinity, inorganic carbon, or base cations (Figures 2 and S4). Organic carbon content was significantly higher in the 20-30 cm depth increment (Figure 3A) and, when summed across the entire column, there was significantly greater total carbon in the MP1-treated soils (Figure 3C). Overall, these data provide evidence of increased weathering (higher pH and soil carbonate alkalinity) and total soil carbon associated with MP1 additions. These findings are similar to those presented in Yang et al. (2026), and despite patterns suggesting a consistent response to the MP1 amendment, the 63-day experiment may not have been long enough relative to the effect size (Hasemer et al., 2024). By further partitioning the soil into pools or fractions via sequential selective dissolution, we improve our ability to understand these suggestive but inconclusive bulk soil data.

4.3 Cation tracing suggests retention of weathering products in secondary minerals


Selective dissolution is a well-established tool for understanding the distribution of elements across a variety of mineralogic, biologic, or geochemical soil fractions in soils and sediments (Heckman et al., 2018; Tessier et al., 1979; Wagai and Mayer, 2007). Although there are some potential pitfalls when using such an approach to compare across different soils (e.g., Hass and Fine, 2010; Martin et al., 1987), when applied to a controlled laboratory experiment, partitioning elements between soil ‘fractions’ or ‘pools’ provides additional information about sources and sinks associated with changes in the overall system chemistry in response to a treatment. In the context of silicate weathering, this information can be further used to understand the potential implications for CDR (Te Pas et al., 2025). The sequential extraction data indicated significant increases in base cations within the carbonate (Figure 4) and reducible (Figure 5) fractions of the MP1-treated columns, with no change in the exchangeable (Figure S5) or oxidizable (Figure S8) fractions.

Given that cation enrichment in carbonate and (hydr)oxides was not accompanied by decreases from any fractions in the MP1-treated columns, remobilization from other depths or fractions is ruled out as a source for the observed increases. Rather, the excess cations, as well as Si and Fe in the reducible fraction, are likely derived from a reservoir not directly quantified with the sequential extraction, with the most likely source being cation-bearing silicate minerals. For example, increases in Fe and Mg in the reducible fraction have been observed following basalt additions (Niron et al., 2024; Vienne et al., 2026).

The depth dependence of the observed carbonate (10-20 cm) and reducible fraction base cation enrichments (20-30 cm) within the MP1-treated columns suggest that microbially enhanced weathering could be spatially variable or that the secondary phases were influenced by the reactive transport and accumulation of weathering products. We did not quantify root abundance with depth, which could have provided insight to the depth distribution of rhizosphere activity. However, because soil respiration generally results in $p\text{CO}_2$ values well above atmospheric levels and the mesocosms are open on the top and bottom, it is reasonable to expect that soil $p\text{CO}_2$ was highest in the intermediate depth. The observed patterns of carbonate and reducible enrichment are consistent with the accumulation of weathering products as porewater infiltrates through the columns. Given that there were no significant differences observed between the treatments in the leachate flux of base cations, it appears that the products of microbially enhanced weathering were largely retained within the soils.

The accumulation of weathering-derived base cations in soil pools, sometimes referred to as 'cation-scavenging,' influences whether or not the associated CDR is realized (Bijma et al., 2026; Hasemer et al., 2024; Steinwider et al., 2025; Te Pas et al., 2025; Vienne et al., 2026). For the purpose of durable CDR, the ideal scenario is that base cations generated from silicate mineral dissolution are exported from the system in the soil porewater, along with the negatively charged carbon alkalinity. Due to the constraint of charge balance in solution, when base cations generated from weathering are retained within the soil, the corresponding alkalinity is also retained, and in some cases (e.g., acidic soils) can be converted back to CO_2 . However, not all soil sinks for base cations are equivalent (Figure 7). The formation of carbonate minerals results in realized CDR, but only half of what would occur if alkalinity was exported. On the other hand, sequestration of base cations in the exchangeable or oxidizable pools are assumed to delay CDR because those cations can eventually be released and exported (Figure 7). Finally, when base cations accumulate in the reducible pool including secondary hydr(oxides) and clays, or are taken up by plants, CDR is inhibited (Steinwider et al., 2025; Te Pas et al., 2025; Vienne et al., 2026).

The molar ratios of Fe, Mg, and Si in the reducible fraction were generally consistent with formation of Si-Mg rich Fe-(hydr)oxides, which could be a precursor to smectite clays. The eventual formation of smectite would be consistent with normal weathering processes in these North Dakota soils (Franzen and Bu, 2023). Although both Fe-(hydr)oxides and smectite tend to have high cation exchange capacity (compared with primary silicates), we did not see significant increases in CEC at any depth in the MP1 treatment (Figure S2). However, we did observe a significant increase in organic carbon in the 20-30 cm increment (Figure 3), which could be related to increased (hydr)oxides as suggested by the reducible fraction data. While more work is needed to definitively identify the minerals associated with the reducible fraction and the possible implications for SOC, it is worth noting, compared with EW approaches where exogenous mineral feedstock are applied to the soil, that the MAW approach is less likely to generate new forms of secondary minerals that were not already present in the soil.



385 Figure 7: The mass balance approach reported here allows for tracking the fate of the observed increases in base cations from the
 acceleration of silicate weathering by *B. subtilis* MP1. For simplification, we assume the observed increase in Ca ions is derived from
 the dissolution of anorthite mineral, and that the resulting Ca ions end up in one of the six possible soil fractions: Carbonate, DIC,
 Exchangeable, Oxidizable, Reducible and Plant. The width of the gray arrows is proportional to the amount of new Ca ions that end
 390 up in one of the six characterized soil fractions. The width of the yellow arrows is proportional to the amount of new Ca ions that
 end up in the exchangeable and carbonate pools, corrected by the soil inorganic carbon measurement. Only Ca ions that end up in
 the carbonate pool or are exported as dissolved inorganic carbon constitute “realized CDR”. The exchangeable and oxidizable pools
 constitute future “potential CDR”, while the reducible and plant pools represent cation scavenging that do not lead to CDR (“no
 CDR”).

395 4.4 The importance of carbonate precipitation in microbially accelerated weathering

The relative increase in base cation content of the carbonate pool in the MP1-treated columns is consistent with increases in soil carbonate reported in Timmermann et al. (2025) and supports for the conceptual model presented above, whereby carbonate precipitation is linked to microbially accelerated silicate weathering (i.e., carbonation). In addition to generating acidity, carbonate formation is a sink for weathering products, maintaining higher rates of weathering by limiting reductions in reaction affinity associated with the accumulation of weathering products (i.e., Maher et al., 2009). In contrast, surface passivation may occur in association with carbonate precipitation, which could slow weathering reactions over longer timescales (Harrison et al., 2015; Power et al., 2021). A more advanced understanding of spatio-temporal partitioning between silicate dissolution and secondary mineral precipitation is necessary to further constrain the role of carbonation in soil weathering systems.

405

It is noteworthy that the sequential extraction results showed a significant increase in the carbonate fraction from 10-20 cm in MP1 treatment, whereas bulk SIC increases in the MP1 treatment were not significant (Figure S2B). This difference could derive from higher variability of SIC compared with the sequential extraction base cation data, or from SIC values approaching analytical detection limits. The SIC measurements are highly correlated with the total base cations in carbonate extraction (Pearson's $r = 0.89$, $p < 0.0001$) but estimates of additional carbonate in the MP1 columns from SIC measurements (0.018 mol CaCO_3 per column) were higher than the sequential extraction base cation estimates (0.010 mol CaCO_3 per column). We attribute this difference to the partial dissolution of newly formed carbonates during the exchangeable cation extraction. Although 1.0 M NH_4OAc is widely used to estimate exchangeable cations, it is known to partially dissolve carbonates under some conditions (Nel et al., 2023; Tessier et al., 1979), potentially leading to an overestimation of exchangeable Ca and Mg and to a corresponding underestimation in the carbonate pool. Furthermore, newly formed biogenic carbonates (e.g., beta calcretes) tend to be fine grained and may be more prone to partial dissolution (Domínguez-Villar et al., 2022; Wright, 1989). Because selective sequential extractions can be susceptible to over- or under-extraction (Hass and Fine, 2010), we assume the SIC-based estimates to be the more accurate measure of the magnitude of soil carbonate.

420 Based on the mass balance calculations, Ca generated from acceleration of silicate weathering was primarily partitioned between the carbonate and exchangeable soil fractions (Figures 6 and 7). However, as discussed above, the sequential extraction method may overestimate the amount of weathering-derived base cations in the exchangeable fraction at the expense of the carbonate fraction. Together these two pools account for the majority of new cations generated through silicate weathering in this soil. The mass of exchangeable Ca (Figure S5) as well as base saturation of the exchange complex (Figure S6) trended higher in the MP1-treated columns but these patterns were not statistically significant, suggesting they may be difficult to detect due to natural soil heterogeneity. Furthermore, in soils with a higher initial base saturation (e.g., Timmermann et al., 2025), we would expect a larger proportion of new Ca to end up in the carbonate fraction. In other words, the base

saturation of the soil will influence the efficiency of the carbonate sink, with higher initial degrees of base saturation likely resulting in more carbonate.

430

There is a large body of evidence demonstrating microbially induced carbonate precipitation by an array of different microbes including *Bacillus subtilis* (Zhu and Ditttrich, 2016). Moreover, we have previously demonstrated that *B. subtilis* MP1 increases the rate of carbonate precipitation in alkaline soils (Timmermann et al., 2025). It follows that carbonate precipitation likely occurs in proximity to the MP1 biofilm, and when compared to EW, microbially accelerated weathering may be more resilient to cation scavenging by soil pools that either delay or inhibit CDR (Figure 7). Additional research is required to better understand the role of biogenic carbonate precipitation in these soil weathering systems.

435

4.5 Cation mass-balance indicates increased silicate weathering in MP1-treated soils

Combining results of the sequential extraction with cation fluxes measured in leachate, and base cations in plant tissue, we found elevated masses of Ca, Mg, and K in the MP1-treated soils, compared to UTC soils (Figure 6). Whereas the total mass of Na decreased in the MP1-treated soils. Taken together, these results provide further support for MP1-driven acceleration of silicate mineral weathering as the ultimate source of the additional cations. On average, the masses of available Ca, Mg, and K in the MP1-treated soils were 6.3%, 2.7%, and 1.1% higher than in the UTC columns, respectively. In contrast, there was a 9.6% reduction of available Na in the MP1-treated columns. However, it should be noted that the magnitudes (Table S3) of MP1-driven changes in Na (-10 mg col⁻¹) and K (+12 mg col⁻¹) were much smaller than Ca (+806 mg col⁻¹) or Mg (+154 mg col⁻¹) and are therefore more likely to be explained by natural heterogeneity or by a limited ability to detect small changes at low concentrations. In total, increases in base cations equate to an average increase of 52.8 mEq of charge per column in the MP1-treated soils, with plant, oxidizable, and leachate fractions accounting for a very small fraction of the total (Figure 6). This corresponds to a maximum CDR of 1.1 g CO₂ kg⁻¹ soil, assuming all cations from increased silicate weathering are exported from the near field as alkalinity. Based on sequential extraction data, the carbonate fraction accounted for 37% (19.5 mEq col⁻¹) of the overall increase, which is very close to estimates for wollastonite weathering (soil pH = 5.16) reported in te Pas et al. (2025). However, when SIC measurements are used to correct for the potential overextraction of cations in the exchangeable step we estimate the carbonate fraction accounts for as much as 35.2 mEq col⁻¹ or 67% of new cations from silicate weathering. Carbonate is a less efficient sink for weathering products than export of alkalinity (Figure 7), still the CDR associated with the carbonate fraction accounts for between 0.20 and 0.36 g CO₂ kg⁻¹ soil and, importantly, it corresponds to realized CDR.

440

445

450

455

4.6 Considerations of the durability of CDR generated from MAW

Under field conditions, the fate of the accumulated carbonate will determine durability of the CDR generated through the MAW approach. If the accumulated carbonates are stable, then the durability of CDR will be at the least equivalent to their soil residence time, which can reach millennia (Landi et al., 2003; Monger et al., 2015). On the other hand, if the accumulated

460

carbonates redissolve, then long-term CDR durability will hinge on whether the associated base cations are exported from the system and transported to the ocean (as with traditional EW systems). Raymond and coauthors (2025) note that carbonate dissolution in buffered soil systems results in the production of bicarbonate (HCO_3^-) rather than CO_2 outgassing. This is supported by field observations from calcareous agricultural soils demonstrating that, even under active fertilization and nitrification, abiotic CO_2 emissions were negligible (Hodges et al., 2021). If, as we suggest, increases in soil carbonate are derived from the microbial acceleration of silicate weathering, any subsequent redissolution of the newly formed carbonates will correspond to an additional mole of CO_2 removed for every mole of CaCO_3 . Furthermore, because of the alkaline pH of the soil, the resulting inorganic carbon will be in the form of HCO_3^- not CO_2 . This implies that an MRV approach for MAW based on the accumulation of SIC represents a conservative estimate of the carbon removal, and the resulting CDR is likely to be highly durable.

5 Conclusions

The results of the controlled mesocosm experiment indicated small but significant differences indicating that the MP1 microbial amendment is associated with increased base cation content, primarily in the secondary carbonate and exchangeable soil fractions, of treated versus control soils. Furthermore, the sequential selective dissolution measurements and cation mass balance calculations provide evidence of silicate weathering increases associated with application of the MP1 microbe and imply that secondary carbonate formation plays an important role in this process. Because there was limited evidence of increased base cation or alkalinity export from the experimental mesocosms, the exchangeable and carbonate pools were the primary sinks for weathering-derived based cations, with smaller but significant increases also observed in the reducible pool.

By combining the soil sequential selective extraction with measurements of soil leachate and plant biomass, we quantified the overall increase in base cations due to silicate weathering and traced the fate of these weathering products (Figure 7). We show that between 37 and 67% of weathering driven increases in base cations went to carbonate formation, with the remaining cations retained mainly in the soil exchangeable pool, which is assumed to delay CDR. At much smaller magnitudes, the remaining cations were retained in the soil organic pool, representing potential future CDR and in the soil reducible pool and plant fraction, which would constitute no CDR. The range reported for carbonate reflects the possible overextraction of the exchangeable pool, which we constrained with independent measurements of SIC. Our findings also suggest caution should be used when comparing sequential extraction results across soils spanning a wide range of pH. In soils with pH greater than ~7, it is also advisable to measure SIC as an independent check on the magnitude of the carbonate fraction.

These results indicate an advantage of using SIC as the basis for measurement, reporting, and verification (MRV) of CDR in neutral to alkaline soils. Although there was a strong correlation between base cation and SIC-based estimates of carbonates, SIC is easier and more cost effective to measure, and in soils with higher pH it may also be more accurate. Furthermore, when

MAW is applied in well-buffered soils, a SIC-based MRV approach represents a conservative estimate of CDR. While field-
495 scale studies and modeling should be used to confirm the long-term durability of CDR associated with MAW, our results
suggest this approach may have significant benefits compared with traditional EW when applied under appropriate conditions.

Data availability

The data described in this work are publicly archived and available (Fuenzalida-Meriz, 2026;
<https://doi.org/10.5281/zenodo.18420180>).

500 Supplement link

The link to the supplement will be included by Copernicus, if applicable.

Author contributions

CRL prepared the manuscript including data analysis, data interpretation, and production of figures. HN conducted the
sequential extraction chemistry and contributed to data analyses. TT designed and oversaw the experiment and contributed to
505 manuscript preparation including writing, editing, and data analyses. PDW conducted chemical analyses, data analyses, and
edited the manuscript. YY ran the mesocosm experiment and conducted biogeochemical analyses. DD conducted laboratory
work and analyses. GFM managed the project and contributed to data analysis, figure production, and edited the manuscript.

Competing interests

Corey R. Lawrence, Tania Timmermann, Philip D. Weyman, Yun-Ya Yang, Daniel Does, and Gonzalo Fuenzalida-Meriz are
510 or were employed by Andes Ag Inc., the company that funded this study.

Disclaimer

Copernicus Publications adds a standard disclaimer: “Copernicus Publications remains neutral with regard to jurisdictional
claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While
Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.
515 Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.”
Please feel free to add disclaimer text at your choice, if applicable.

Acknowledgements

The authors thank Dr. Sara Vicca (University of Antwerp) who contributed to early discussions of the selective sequential extraction procedure and provided the laboratory facilities for that work. /

520 Review statement

The review statement will be added by Copernicus Publications listing the handling editor as well as all contributing referees according to their status anonymous or identified.

References

- Banfield, J. F., Barker, W. W., Welch, S. A., and Taunton, A.: Biological impact on mineral dissolution: Application of the
525 lichen model to understanding mineral weathering in the rhizosphere, *Proc. Natl. Acad. Sci.*, 96, 3404–3411,
<https://doi.org/10.1073/pnas.96.7.3404>, 1999.
- Berner, R. A.: The Rise of Plants and Their Effect on Weathering and Atmospheric CO₂, *Science*, 276, 544–546,
<https://doi.org/10.1126/science.276.5312.544>, 1997.
- Bijma, J., Hagens, M., Hammes, J. S., Planavsky, N., Pogge Von Strandmann, P. A. E., Reershemius, T., Reinhard, C. T.,
530 Renforth, P., Suhrhoff, T. J., Vicca, S., Vienne, A., and Wolf-Gladrow, D.: Reviews and syntheses: Carbon vs. cation based
MRV of Enhanced Rock Weathering and the issue of soil organic carbon, *Biogeosciences*, 23, 53–75,
<https://doi.org/10.5194/bg-23-53-2026>, 2026.
- Clarkson, M. O., Larkin, C. S., Swoboda, P., Reershemius, T., Suhrhoff, T. J., Maesano, C. N., and Campbell, J. S.: A review
of measurement for quantification of carbon dioxide removal by enhanced weathering in soil, *Front. Clim.*, 6, 1345224,
535 <https://doi.org/10.3389/fclim.2024.1345224>, 2024.
- Corbett, T., Odelius, E., Uebel, T., Jonsson, T., Singh, A., Vicca, S., Hagens, M., Budnyak, T., Tkachenko, O., Poetra, R.,
Hartmann, J., Janssens, I. A., Niron, H., Van Tendeloo, M., Vlaeminck, S. E., and Neubeck, A.: Microbial dissolution of Gran
Canaria lapilli in small-scale flow through columns: carbon dioxide removal potential, *Npj Mater. Degrad.*, 9, 60,
<https://doi.org/10.1038/s41529-025-00611-9>, 2025.
- 540 Domínguez-Villar, D., Bensa, A., Švob, M., and Krklec, K.: Causes and implications of the seasonal dissolution and
precipitation of pedogenic carbonates in soils of karst regions – A thermodynamic model approach, *Geoderma*, 423, 115962,
<https://doi.org/10.1016/j.geoderma.2022.115962>, 2022.
- Dong, H., Huang, L., Zhao, L., Zeng, Q., Liu, X., Sheng, Y., Shi, L., Wu, G., Jiang, H., Li, F., Zhang, L., Guo, D., Li, G., Hou,
W., and Chen, H.: A critical review of mineral–microbe interaction and co-evolution: mechanisms and applications, *Natl. Sci.*
545 *Rev.*, 9, nwac128, <https://doi.org/10.1093/nsr/nwac128>, 2022.

- Fierer, N. and Walsh, C. M.: Can we manipulate the soil microbiome to promote carbon sequestration in croplands?, *PLOS Biol.*, 21, e3002207, <https://doi.org/10.1371/journal.pbio.3002207>, 2023.
- Finlay, R. D., Mahmood, S., Rosenstock, N., Bolou-Bi, E. B., Köhler, S. J., Fahad, Z., Rosling, A., Wallander, H., Belyazid, S., Bishop, K., and Lian, B.: Reviews and syntheses: Biological weathering and its consequences at different spatial levels –
 550 from nanoscale to global scale, *Biogeosciences*, 17, 1507–1533, <https://doi.org/10.5194/bg-17-1507-2020>, 2020.
- Franzen, D. W. and Bu, H.: North Dakota Clay Mineralogy Impacts Crop Potassium Nutrition and Tillage Systems, North Dakota State University, Fargo, ND, USA, 2023.
- Fuenzalida-Meriz, G. A.: Dataset for manuscript “Microbially-accelerated weathering achieves carbon dioxide removal by coupling silicate dissolution with carbonate precipitation,” <https://doi.org/10.5281/zenodo.18420180>, 2026.
- 555 Harrison, A. L., Dipple, G. M., Power, I. M., and Mayer, K. U.: Influence of surface passivation and water content on mineral reactions in unsaturated porous media: Implications for brucite carbonation and CO₂ sequestration, *Geochim. Cosmochim. Acta*, 148, 477–495, <https://doi.org/10.1016/j.gca.2014.10.020>, 2015.
- Hasemer, H., Borevitz, J., and Buss, W.: Measuring enhanced weathering: inorganic carbon-based approaches may be required to complement cation-based approaches, *Front. Clim.*, 6, 1352825, <https://doi.org/10.3389/fclim.2024.1352825>, 2024.
- 560 Hass, A. and Fine, P.: Sequential Selective Extraction Procedures for the Study of Heavy Metals in Soils, Sediments, and Waste Materials—a Critical Review, *Crit. Rev. Environ. Sci. Technol.*, 40, 365–399, <https://doi.org/10.1080/10643380802377992>, 2010.
- Heckman, K., Lawrence, C. R., and Harden, J. W.: A sequential selective dissolution method to quantify storage and stability of organic carbon associated with Al and Fe hydroxide phases, *Geoderma*, 312, 24–35,
 565 <https://doi.org/10.1016/j.geoderma.2017.09.043>, 2018.
- Hodges, C., Brantley, S. L., Sharifironizi, M., Forsythe, B., Tang, Q., Carpenter, N., and Kaye, J.: Soil Carbon Dioxide Flux Partitioning in a Calcareous Watershed With Agricultural Impacts, *J. Geophys. Res. Biogeosciences*, 126, e2021JG006379, <https://doi.org/10.1029/2021JG006379>, 2021.
- Hopf, J., Langenhorst, F., Pollok, K., Merten, D., and Kothe, E.: Influence of microorganisms on biotite dissolution: An
 570 experimental approach, *Geochemistry*, 69, 45–56, <https://doi.org/10.1016/j.chemer.2008.11.001>, 2009.
- Johnson, D. M., Hooper, P. R., and Conrey, R. M.: XRF Analysis of Rocks and Minerals for Major and Trace Elements on a Single Low Dilution Li-tetraborate Fused Bead, 1999.
- Johnstone, I., Fuss, S., Walsh, N., and Höglund, R.: Carbon markets for carbon dioxide removal, *Clim. Policy*, 26, 151–158, <https://doi.org/10.1080/14693062.2025.2478288>, 2026.
- 575 Kelly, E. F., Chadwick, O. A., and Hilinski, T. E.: The Effect of Plants on Mineral Weathering, *Biogeochemistry*, 42, 21–53, <https://doi.org/10.1023/A:1005919306687>, 1998.
- Landi, A., Mermut, A. R., and Anderson, D. W.: Origin and rate of pedogenic carbonate accumulation in Saskatchewan soils, Canada, *Geoderma*, 117, 143–156, [https://doi.org/10.1016/S0016-7061\(03\)00161-7](https://doi.org/10.1016/S0016-7061(03)00161-7), 2003.

- 580 Maher, K., Steefel, C. I., White, A. F., and Stonestrom, D. A.: The role of reaction affinity and secondary minerals in regulating chemical weathering rates at the Santa Cruz Soil Chronosequence, California, *Geochim. Cosmochim. Acta*, 73, 2804–2831, <https://doi.org/10.1016/j.gca.2009.01.030>, 2009.
- Martin, J. M., Nirel, P., and Thomas, A. J.: Sequential extraction techniques: Promises and problems, *Mar. Chem.*, 22, 313–341, [https://doi.org/10.1016/0304-4203\(87\)90017-X](https://doi.org/10.1016/0304-4203(87)90017-X), 1987.
- 585 McConnaughey, T. A. and Whelan, J. F.: Calcification generates protons for nutrient and bicarbonate uptake, *Earth-Sci. Rev.*, 42, 95–117, [https://doi.org/10.1016/S0012-8252\(96\)00036-0](https://doi.org/10.1016/S0012-8252(96)00036-0), 1997.
- Monger, H. C., Kraimer, R. A., Khresat, S., Cole, D. R., Wang, X., and Wang, J.: Sequestration of inorganic carbon in soil and groundwater, *Geology*, 43, 375–378, <https://doi.org/10.1130/G36449.1>, 2015.
- Nel, T., Bruneel, Y., and Smolders, E.: Comparison of five methods to determine the cation exchange capacity of soil, *J. Plant Nutr. Soil Sci.*, 186, 311–320, <https://doi.org/10.1002/jpln.202200378>, 2023.
- 590 Nelson, D. W. and Sommers, L. E.: Total Carbon, Organic Carbon, and Organic Matter, in: *Methods of Soil Analysis*, John Wiley & Sons, Ltd, 961–1010, <https://doi.org/10.2136/sssabookser5.3.c34>, 1996.
- Niron, H., Vienne, A., Frings, P., Poetra, R., and Vicca, S.: Exploring the synergy of enhanced weathering and *Bacillus subtilis*: A promising strategy for sustainable agriculture, *Glob. Change Biol.*, 30, e17511, <https://doi.org/10.1111/gcb.17511>, 2024.
- 595 Niron, H., Calogiuri, T., Singh, A., Neubeck, A., Van Tendeloo, M., Cox, T., Hartmann, J., Vlaeminck, S. E., and Vicca, S.: Alkalinity production and carbon capture from dunite weathering: Individual effects of oxalate, citrate, and EDTA salts, *Chem. Eng. J. Adv.*, 24, 100902, <https://doi.org/10.1016/j.cej.2025.100902>, 2025.
- Peters, J., Nathan, M., Lakoski, C., and Nathan, M.: Chapter 4: pH and Lime Requirement, in: *Recommended Chemical Soil Test Procedures for the North Central Region*, vol. 221, University of Missouri, 16–22, 2012.
- 600 Power, I. M., Paulo, C., Long, H., Lockhart, J. A., Stubbs, A. R., French, D., and Caldwell, R.: Carbonation, Cementation, and Stabilization of Ultramafic Mine Tailings, *Environ. Sci. Technol.*, 55, 10056–10066, <https://doi.org/10.1021/acs.est.1c01570>, 2021.
- Raymond, P., Planavsky, N., and Reinhard, C. T.: Using carbonates for carbon removal, *Nat. Water*, 3, 844–847, <https://doi.org/10.1038/s44221-025-00473-0>, 2025.
- 605 Ribeiro, I. D. A., Volpiano, C. G., Vargas, L. K., Granada, C. E., Lisboa, B. B., and Passaglia, L. M. P.: Use of Mineral Weathering Bacteria to Enhance Nutrient Availability in Crops: A Review, *Front. Plant Sci.*, 11, 590774, <https://doi.org/10.3389/fpls.2020.590774>, 2020.
- Richards, L. A., Allison, L. E., Bernstein, L., Bower, C. A., Brown, J. W., Fireman, M., Hatcher, J. T., Hayward, H. E., Pearson, G. A., and Reeve, R. C.: *Diagnosis and Improvement of Saline and Alkali Soils*, edited by: Richards, L. A., Soil and Water Conservation Research Branch Agricultural Research Service, 1954.
- 610

- Samuels, T., Bryce, C., Landenmark, H., Marie-Loudon, C., Nicholson, N., Stevens, A. H., and Cockell, C.: Microbial Weathering of Minerals and Rocks in Natural Environments, in: Biogeochemical Cycles, American Geophysical Union (AGU), 59–79, <https://doi.org/10.1002/9781119413332.ch3>, 2020.
- Schulte, E. E. and Hopkins, B. G.: Estimation of Soil Organic Matter by Weight Loss-On-Ignition, in: Soil Organic Matter: Analysis and Interpretation, John Wiley & Sons, Ltd, 21–31, <https://doi.org/10.2136/sssaspecpub46.c3>, 1996.
- Smith, S., Geden, O., Gidden, M., Lamb, W., Nemet, G., Minx, J., Buck, H., Burke, J., Cox, E., and Edwards, M.: The State of Carbon Dioxide Removal, 2nd Edition, OSF, <https://doi.org/10.17605/OSF.IO/F85QJ>, 2024.
- Song, W., Ogawa, N., Oguchi, C. T., Hatta, T., and Matsukura, Y.: Effect of *Bacillus subtilis* on granite weathering: A laboratory experiment, CATENA, 70, 275–281, <https://doi.org/10.1016/j.catena.2006.09.003>, 2007.
- Steinwider, L., Boito, L., Frings, P. J., Niron, H., Rijnders, J., De Schutter, A., Vienne, A., and Vicca, S.: Beyond Inorganic C: Soil Organic C as a Key Pathway for Carbon Sequestration in Enhanced Weathering, Glob. Change Biol., 31, e70340, <https://doi.org/10.1111/gcb.70340>, 2025.
- Steinwider, L., Boito, L., de Schutter, A., Frings, P. J., Miladinović, N., Niron, H., Rijnders, J., Roussard, J., van Acker, K., van Gerven, T., Vienne, A., Watjanatepin, P., and Vicca, S.: Higher Inorganic CO₂ Removal Despite Slower Weathering in an Enhanced Weathering Experiment With Steel Slags and Basalt, Glob. Change Biol., 32, e70666, <https://doi.org/10.1111/gcb.70666>, 2026.
- Sumner, M. E. and Miller, W. P.: Cation Exchange Capacity and Exchange Coefficients, in: Methods of Soil Analysis, Part 3: Chemical Methods, vol. 5, edited by: Sparks, D. L., Page, A. L., Helmke, P. A., and Loeppert, R. H., American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, 1201–1229, 2020.
- Supuran, C. T.: Structure and function of carbonic anhydrases, Biochem. J., 473, 2023–2032, <https://doi.org/10.1042/BCJ20160115>, 2016.
- Te Pas, E. E. E. M., Chang, E., Marklein, A. R., Comans, R. N. J., and Hagens, M.: Accounting for retarded weathering products in comparing methods for quantifying carbon dioxide removal in a short-term enhanced weathering study, Front. Clim., 6, 1524998, <https://doi.org/10.3389/fclim.2024.1524998>, 2025.
- Tessier, A., Campbell, P. G. C., and Bisson, M.: Sequential extraction procedure for the speciation of particulate trace metals, Anal. Chem., 51, 844–851, <https://doi.org/10.1021/ac50043a017>, 1979.
- Timmermann, T., Yip, C., Yang, Y., Wemmer, K. A., Chowdhury, A., Dores, D., Takayama, T., Nademanee, S., Traag, B. A., Zamanian, K., González, B., Breecker, D. O., Fierer, N., Slessarev, E. W., and Fuenzalida-Meriz, G. A.: Harnessing Microbes to Weather Native Silicates in Agricultural Soils for Scalable Carbon Dioxide Removal, Glob. Change Biol., 31, e70216, <https://doi.org/10.1111/gcb.70216>, 2025.
- Tosca, N. J. and Tutolo, B. M.: Alkalinity in Theory and Practice, Elements, 19, 7–9, <https://doi.org/10.2138/gselements.19.1.7>, 2023.
- Uhlig, D. and Von Blanckenburg, F.: How Slow Rock Weathering Balances Nutrient Loss During Fast Forest Floor Turnover in Montane, Temperate Forest Ecosystems, Front. Earth Sci., 7, 159, <https://doi.org/10.3389/feart.2019.00159>, 2019.

- 645 Verbruggen, E., Struyf, E., and Vicca, S.: Can arbuscular mycorrhizal fungi speed up carbon sequestration by enhanced weathering?, *PLANTS PEOPLE PLANET*, 3, 445–453, <https://doi.org/10.1002/ppp3.10179>, 2021.
- Vicca, S., Goll, D. S., Hagens, M., Hartmann, J., Janssens, I. A., Neubeck, A., Peñuelas, J., Poblador, S., Rijnders, J., Sardans, J., Struyf, E., Swoboda, P., van Groenigen, J. W., Vienne, A., and Verbruggen, E.: Is the climate change mitigation effect of enhanced silicate weathering governed by biological processes?, *Glob. Change Biol.*, 28, 711–726, <https://doi.org/10.1111/gcb.15993>, 2022.
- 650 Vienne, A., Frings, P., Rijnders, J., Boito, L., Hartmann, J., Niron, H., Poetra, R., Estrada, M. P., Reershemius, T., Steinwigger, L., Suhrhoff, T. J., and Vicca, S.: Weathering without realizing inorganic CO₂ removal revealed through base cation monitoring, *SOIL*, 12, 421–440, <https://doi.org/10.5194/soil-12-421-2026>, 2026.
- Wagai, R. and Mayer, L. M.: Sorptive stabilization of organic matter in soils by hydrous iron oxides, *Geochim. Cosmochim. Acta*, 71, 25–35, <https://doi.org/10.1016/j.gca.2006.08.047>, 2007.
- 655 Walker, J. C. G., Hays, P. B., and Kasting, J. F.: A negative feedback mechanism for the long-term stabilization of Earth’s surface temperature, *J. Geophys. Res. Oceans*, 86, 9776–9782, <https://doi.org/10.1029/JC086iC10p09776>, 1981.
- Wang, J., Qiu, M., Shen, Z., Chen, L., and Ge, Y.: Microbial modulation of soil pH: A self-benefiting mechanism exemplified by *Bacillus*, *Soil Biol. Biochem.*, 210, 109949, <https://doi.org/10.1016/j.soilbio.2025.109949>, 2025.
- 660 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemond, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H.: Welcome to the Tidyverse, *J. Open Source Softw.*, 4, 1686, <https://doi.org/10.21105/joss.01686>, 2019.
- Wild, B., Gerrits, R., and Bonneville, S.: The contribution of living organisms to rock weathering in the critical zone, *Npj Mater. Degrad.*, 6, 98, <https://doi.org/10.1038/s41529-022-00312-7>, 2022.
- 665 Wright, V. P.: Terrestrial stromatolites and laminar calcretes: a review, *Sediment. Geol.*, 65, 1–13, [https://doi.org/10.1016/0037-0738\(89\)90002-X](https://doi.org/10.1016/0037-0738(89)90002-X), 1989.
- Xiao, L., Lian, B., Hao, J., Liu, C., and Wang, S.: Effect of carbonic anhydrase on silicate weathering and carbonate formation at present day CO₂ concentrations compared to primordial values, *Sci. Rep.*, 5, 7733, <https://doi.org/10.1038/srep07733>, 2015.
- 670 Yang, Y., Bueno De Mesquita, C. P., Lawrence, C. R., Weyman, P. D., Dore, D., Timmermann, T., Fierer, N., and Fuenzalida-Meriz, G. A.: Synergistic Effects of a Microbial Amendment and Crushed Basalt: Soil Geochemical and Microbial Responses, *Glob. Change Biol.*, 32, e70705, <https://doi.org/10.1111/gcb.70705>, 2026.
- Yip, C., Weyman, P. D., Wemmer, K. A., Yang, Y.-Y., Chowdhury, A., Traag, B. A., Timmermann, T., and Fuenzalida-Meriz, G.: Quantification of soil inorganic carbon using sulfamic acid and gas chromatography, *PLOS One*, 20, e0320778, <https://doi.org/10.1371/journal.pone.0320778>, 2025.
- 675 Zhu, T. and Dittrich, M.: Carbonate Precipitation through Microbial Activities in Natural Environment, and Their Potential in Biotechnology: A Review, *Front. Bioeng. Biotechnol.*, 4, <https://doi.org/10.3389/fbioe.2016.00004>, 2016.